

# ZytoDot® 2C SPEC MET/CEN 7 Probe

CE  
0124  
IVD


## Background

The ZytoDot® 2C SPEC MET/CEN 7 Probe (PD37) is intended to be used for the qualitative detection of amplifications involving the human MET gene as well as the detection of chromosome 7 alpha satellites in formalin-fixed, paraffin-embedded specimens by chromogenic *in situ* hybridization (CISH). The probe is intended to be used in combination with the ZytoDot® 2C CISH Implementation Kit (Prod. No. C-3044-10/-40). The product is intended for professional use only. All tests using the product should be performed in a certified, licensed anatomic pathology laboratory under the supervision of a pathologist/human geneticist by qualified personnel. The probe is intended to be used as an aid to the differential diagnosis of various cancers and therapeutic measures should not be initiated based on the test result alone.


## Probe Description

The ZytoDot® 2C SPEC MET/CEN 7 Probe is composed of:

- Digoxigenin-labeled polynucleotides (~1.1 ng/μl), which target sequences mapping in 7q31.2\*\* (chr7:116,298,989-116,718,699) harboring the MET gene region.
- Dinitrophenyl-labeled polynucleotides (~1.1 ng/μl), which target sequences mapping in 7p11.1-q11.1 specific for the alpha satellite centromeric region D7Z1 of chromosome 7.
- Formamide based hybridization buffer




Ideogram of chromosome 7 indicating the hybridization locations.



SPEC MET Probe map (not to scale).

## Results

In a normal interphase nucleus, using the ZytoDot® 2C CISH Implementation Kit two red (CEN 7) and two green (MET) signals are expected. In a cell with amplification of the MET gene locus, multiple copies of the green signal or green signal clusters will be observed.



Example of an aberrant signal pattern: Lung cancer tissue section with multiple copies of chromosome 7 (red) and extra MET signals (green) in the nuclei.

Prod. No.	Product	Label	Tests* (Volume)
C-3057-400	ZytoDot 2C SPEC MET/CEN 7 Probe CE 0124 IVD	DIG/DNP	40 (400 μl)
<b>Related Products</b>			
C-3044-40	ZytoDot 2C CISH Implementation Kit CE IVD		40
Incl. Heat Pretreatment Solution EDTA, 500 ml; Pepsin Solution, 4 ml; Wash Buffer SSC, 560 ml; 20x Wash Buffer TBS, 2x 50 ml; Anti-DIG/DNP-Mix, 4 ml; HRP/AP-Polymer-Mix, 4 ml; AP-Red Solution A, 0.4 ml; AP-Red Solution B, 15 ml; HRP-Green Solution A, 0.8 ml; HRP-Green Solution B, 15 ml; Nuclear Blue Solution, 20 ml; Mounting Solution (alcoholic), 4 ml			

\* Using 10 μl probe solution per test. IVD labeled products are only available in certain countries. All other countries research use only! Please contact your local dealer for more information.

\*\*According to Human Genome Assembly GRCh37/hg19